arXiv:astro-ph/9911437v1 23 Nov 1999

Constraints on the Spatial Distribution and Luminosity function of GRBs

Tomasz Bulik

Nicolaus Copernicus Astronomical Center
Bartycka 18, 00716 Warsaw, Poland

Abstract. The distance scale to gamma ray bursts has been a subject of scientific debate for almost thirty years. Up to the discovery of afterglows only indirect methods could be used to constrain the distance scale to this objects. I review some of these results, and show the current limits on the spatial distribution and luminosity function of GRBs. The results obtained with different methods indicate that gamma-ray bursts lie at the typical redshifts z_{ave} between 1 and 2, however there can exist a long tail of the distribution stretching to higher redshifts. The width of the GRB luminosity function (the ratio of the intrinsic brightness of the brightest to the dimmest observed burst) estimated from Beppo SAX bursts with redshifts is almost 10^3.

1. Introduction

Ever since the discovery of gamma ray bursts Klebesadel et al. (1973) the main questions posed by the astronomers in this field were what is the distance scale to these phenomena, and consequently what is their spatial distribution. For almost three decades the main tools for probing the distance scale to gamma-ray bursts were studies of their statistical properties, and model dependent physical arguments. In fact in the Great Debate in 1995 the main argument for the cosmological distance scale was the isotropy of GRBs on the sky (Paczynski, 1995; Lamb, 1995). Apart from the distribution studies there have been a number of attempts to measure the distance scale, and the spatial distribution of GRBs using different methods, based on e.g. the temporal or spectral studies, searches for gravitational lensing etc. Since the discovery of afterglows (Costa et al., 1997b) and measurement of GRB redshifts Metzger et al. (1997a) we have entered a new era in the GRB research and now we can probe directly the spatial distribution of GRBs. The paper is organized as follows: in section 2 I analyze the angular distribution of GRBs, in section 3 I review the constraints on their spatial distribution, in section 4 classes of GRBs are discussed, and the results are summarized in section 5.

2. Angular distribution

The angular distribution of gamma ray bursts is very close to isotropy. In the current BATSE catalogue (Meegan et al., 1998; Paciesas et al., 1999) we



Figure 1. Cumulative distributions of the celestial coordinates of GRBs in the public archive; latitude b on the left, and longitude θ on the right. The thin lines are the distribution expected in the case of isotropy, and the thick lines are the actual observations.

find that the galactic dipole and quadrupole moments are consistent with the values expected for isotropy when taking into account the nonuniform exposure. The current measured value of the dipole moments in the galactic coordinates is $\langle \cos\theta \rangle = -0.024 \pm 0.014$, with the expected value equal to $\langle \cos\theta \rangle_{exp} = -.009 \pm 0.002$, while the quadrupole moment is $\langle \sin^2 b - 1/3 \rangle = 0.0005 \pm 0.0074$, compared to the expected value $\langle \sin^2 b - 1/3 \rangle_{exp} = 0.004 \pm 0.002$.

While the integral moments do not describe the entire distribution, one can perform a more detailed comparison using the Kolmogorov-Smirnov (KS) tests of the distribution of the galactic coordinates of the bursts. The KS test comparison shows an excellent agreement between the expected and observed distributions of right ascensions α, where the probability that the distributions are identical is 97%; the probability for the distribution of declinations is smaller - 22%. This is probably due to uncertainty of the BATSE sky exposure map. This uncertainty is also seen when comparing the galactic coordinates distribution: here the KS probabilities are 14% for galactic latitude distribution and 32% for the galactic longitude distribution.

Thus the distribution of GRBs is consistent with isotropy on the sky. In the first BATSE catalogue (Fishman et al., 1994) there was an evidence of variation of the galactic moments with brightness (Quashnock and Lamb, 1993a). This effect disappeared in the later catalogues. Therefore in the remaining part of the paper I assume that the spatial distribution of GRBs can be factorized into the independent angular and radial parts, and that the angular part is isotropic. The most comprehensive studies of the isotropy and comparisons with anisotropic models have been performed by Loredo and Wasserman (1995, 1998b).



Figure 2. Cumulative distributions of the galactic coordinates of GRBs in the public archive; latitude b on the left, and longitude θ on the right.

3. Spatial distribution

Various statistical methods were used for analyzing the gamma-ray burst brightness distributions. Some authors divided the observed bursts in bins and used a χ^2 statistics to compare the number of bursts in these bins with models. The shortcoming of this method is that the results depend on the arbitrary parameter (or parameters), i.e. the number and the width of the bins. This makes the calculation of statistical significance difficult, since in most cases the authors do not discuss the effects of the bin width. Usually the bins are chosen in such a way to ensure that the number of counts in each bin allows using the Gaussian approximation. Another possibility is to use the Kolmogorov Smirnov test, which measures the largest deviation between the model and the expected cumulative distribution. It seems that the most sensitive, yet also most computationally demanding method is the maximum likelihood method, which takes into account all information contained in the data.

3.1. Brightness distribution

Studies of GRB brightness distribution require a definition of brightness of a given burst. Several measures of brightness have been used: e.g. the peak flux, or the fluence. Each measure of brightness introduces systematic instrumental effects. The measured fluence depends on the background level in a particular detector and on the true time history of a burst. Given that the fluence of a given burst will not necessarily follow the r^{-2} law, even for simple, one peak bursts. The peak flux suffers from the so called Meegan bias. Given a burst with a couple peaks we will tend to choose the moment of the maximum flux as the one where the Poisson fluctuation upwards were the largest. This effect will lead to overestimates of the flux, and the amount of change will depend on the particular time history of the burst. The peak flux distribution will also depend on the particular timescale on which this peak flux was measured. The value of



Figure 3. Brightness distributions of BATSE GRBs. The left panel shows the distributions of peak fluxes on the following time scales from left to right 64ms (thin line), 256ms (medium line), and 1024 ms (thick line). The right panel shows the distribution of peak fluxes: summed over the BATSE channels (solid line) and for channels 2 and 3 (dashed line).

each brightness measure depends also on the spectral response of the detector, and on the spectral interval considered. The distributions of peak fluxes, and fluences of GRBs are shown in Figure 3.

The observed bursting rate R of GRBs can be described by

$$\frac{dR}{dz\, d\Phi\, dL} = \frac{n(z)}{1+z} 4\pi r_z^2 \frac{dr_z}{dz} f(L)\delta(\Phi - \Phi_{det}) \qquad (1)$$

where $n(z)$ is the density of the sources, r_z is the luminosity distance, Φ is the flux, and $f(L)$ is the luminosity function. In general we measure only the observed flux Φ, however, recently for some burst also the redshifts z have been measured. Moreover, in a detailed calculation K-corrections should be added to equation 1.

The simplest model of cosmological GRBs is with standard candle luminosity function, and no source evolution. Such a model has been considered by a number of authors; Mao and Paczynski (1992) showed using an V/V_{max} that the brightness distribution of GRBs is consistent with a simple cosmological interpretation with no free parameters. They found that the maximum redshift is constrained to be $1.0 < z_{max} < 2.5$; see also (Dermer, 1992; Piran, 1992; Dermer, 1992). Wickramasinghe et al. (1993) found $0.9 < z_{max} < 2.0$ using the C_{max}/C_{min} comparison. Cohen and Piran (1995) used the KS-test of the number count distribution in the 2B catalogue and found that the maximal redshift must be $1.4 < z_{max} < 3.1$. In a detailed study with the use of maximum likelihood analysis of the 3B data Loredo and Wasserman (1998a) constrained the standard candle luminosity of GRBs to be $L \approx 0.74 \times 10^{51} \mathrm{erg\,s}^{-1}$.

The effects of the GRB rate density evolution are usually included by assuming that $n(z) \propto (1+z)^D$. In general the results show that the higher the

exponent D, the more luminous GRBs are. The luminosity of a standard candle bursts grows very rapidly with the exponent D, however the details vary from one paper to another and depend on the particular data set used (BATSE or BATSE and PVO) (Cohen and Piran, 1995; Meszaros and Meszaros, 1996; Horvath et al., 1996; Loredo and Wasserman, 1998a)

Luminosity function When considering the luminosity function of GRBs one has to distinguish between the observed and intrinsic luminosity functions. The intrinsic luminosity function is the distribution of luminosities of all bursts regardless of whether we see them or not, while the observed luminosity function is the distribution of luminosities of the observed bursts only. It has been shown that the two distributions are different, and may even have different slopes. The luminosity function is usually represented by a power on a fixed interval, however other functional forms like e.g. the lognormal distribution are also used.

There were several papers discussing the effects of the luminosity function on the shape of the brightness distribution. Horack et al. (1994) used the integral moment analysis and concluded that the width of the observed luminosity function must be smaller than a factor of 6.5. Cohen and Piran (1995) used a two δ functions approximation of the luminosity function and constrained the width of such luminosity function to less than a factor of 14 in the case of BATSE data and less than a factor of 2 for the combined BATSE and PVO data sets. Woods and Loeb (1995) parameterized the luminosity function as a lognormal distribution, used KS-test to compare with the observed C_{max} distribution, and found that the width of the luminosity function must be less than about a factor of 40. Some studies of the effects of the luminosity function were done assuming a halo-core like spatial distribution: Ulmer and Wijers (1995) modeled the BATSE brightness distribution using the maximum likelihood method and found essentially no limits on the intrinsic luminosity function, however Ulmer et al. (1995) used the combined BATSE+PVO) dataset and found the the observed luminosity function width is less than 10. Bulik et al. (1998) also considered a halo-like distribution and using the KS-test found that BATSE data is consistent with the luminosity function wider than a factor of 100. Analyses of the combined BATSE+PVO dataset with χ^2 yielded constraints that the luminosity function width must be less than a factor of 1000 (Horvath et al., 1996; Hakkila et al., 1996). Loredo and Wasserman (1998a) analyze the gamma ray burst luminosity function using Bayesian statistics and maximum likelihood method. They find that the current data set (BATSE 3B catalogue) does not constrain the intrinsic luminosity function. The main reason is that the differences between the models should show only for much fainter fluxes than the BATSE threshold (see Fig. 25 in their paper) . They also find the width of the *observed* luminosity function is around 10^4.

Thus the studies of the GRB luminosity function by analysis of the brightness distribution has taken an unusual direction: initially with small amount of data it seemed that the width of the luminosity function is small, yet with the increasing amount of data and improved statistical methods it appeared that its width can actually be large. There are no strong constraints on the width of the intrinsic luminosity function of GRBs from the brightness distribution.



Figure 4. The left panel shows the distribution of BATSE measured burst durations T_{50} and T_{90}. While a clear correlation between these two measures exists there is also a significant spread. The right panel shows the BATSE bursts in the plane spanned by T_{90} and the peak flux on the 256ms timescale. No clear correlation exists, which shows how difficult it is to find the burst distance scale from brightness duration correlation.

3.2. Temporal studies

It is natural to attempt to use the temporal characteristic of GRBs to find constraints on the distance scale within the framework of the cosmological model. Assuming that weaker bursts are typically further away they should also be longer, due to cosmological redshift. It should be noted that it is difficult to obtain an absolute scale without a physical model or knowledge of a clock in bursts. One can only measure the relative distance (redshift) between groups of weak and bright bursts. Thus the absolute distance scale can be obtained then assuming a distance to the weakest bursts. The method is based on finding a correlation between brightness and duration, or other measure of the intrinsic timescale in gamma-ray burst.

Studies of temporal characteristics of GRBs require defining a fundamental timescale for GRBs. This is not easy, since GRB time histories are very diverse. BATSE catalogue provides two measures of duration T_{50}, and T_{90}, the time in which 50 and 90 percent of the photons arrived. I present the correlation between these two measures of duration in left panel of Figure 4. The spread in these correlation is significant and this shows that there are possible systematic effects that may affect the brightness - duration studies. The right panel of Figure 4 shows the distribution of BATSE GRBs in the duration (T_{90}) and brightness (peak flux on the 256 ms timescale) plane. No correlation is apparent in this graph. Thus several authors decided to analyze the BATSE data themselves and measure durations, or time scales defined differently.

We present a sample of the results obtained with the use of different methods, different measurements of burst durations in Table 1. Typically regardless of the method a stretch factor of approximately 2 was found, meaning that the

Table 1. GRB time stretching results.

Method	S	reference
Peak align	2.25	Norris et al. (1994)
Fluence/Peak	2.0	Norris et al. (1994)
Wavelets	2.25	Norris et al. (1994)
Pulse width	1.8	Davis (1995)
Counts in peak	2.2	Davis (1995)
Total counts	2.0	Davis (1995)
Duration	2.2	Norris et al. (1995)
Autocorrelation	2.0	Fenimore et al. (1995)
Ave. time hist.	1.0	Mitrofanov et al. (1996)
Stretch exponential	2.0	Stern (1996)
Fluence vs T_{50}	1.0	Lee and Petrosian (1997)

typical clock for weak bursts runs slower than that for the bright ones. There are some exceptions: Mitrofanov et al. (1996) and Lee and Petrosian (1997) found no difference between the bright and the dim bursts. The stretch factor of ≈ 2 was interpreted as the redshift of the dim burst of approximately $z_{dim} = 2$. However, Fenimore and Bloom (1995) noticed that in each given burst the width of the peaks is larger when measuring low energy channels than in the case of high energy ones. This leads to the so-called W-correction; the effects of redshift is not only slowing down the clocks but also shifting to lower energies, where the peaks are wider. The inclusion of this correction lead to a new limit on the redshift of the dim bursts $z_{dim} \approx 6$ for the stretch factor of 2. It has to be noted that it is always difficult to separate the effects of source evolution from the true cosmological effect in these studies. In fact Stern et al. (1997) interpreted the variation in the slope of decay of GRB profiles as due to the evolution of sources.

3.3. Host galaxy limits

The host galaxy problem, for quite a while called the "no host problem" was first presented by Schaeffer (1992). He analyzed the contents of error boxes of brightest bursts with the accurate IPN localization, in search of galaxies and found that these boxes contain no galaxies to the limit of twentieth magnitude, while expecting to find nearby galaxies of magnitude ≈ 16 in them. Of course, in this calculation it was assumed that GRBs are not totally exotic phenomena and therefore are taking place in galaxies. This finding led to the conclusion that the standard cosmological model (with typical GRB luminosity 6×10^{50}ergs) has some difficulty. An analysis of the infrared galaxy catalog by Larson and MCLean (1997), was claimed to show consistency of the contents of GRB errors boxes with the cosmological model, however Band et al. (1999) showed using Bayesian analysis that the infrared data set provided no really useful limits.

Recently this problem has been revisited by Schaeffer (1998) and Band et al. (1999). Schaeffer (1998) analyzed the contents of 26 small GRB error boxes and concluded that the lower limit on GRB luminosity is $6 \times 10^{58}\,\mathrm{phot\,cm^2\,s^{-1}}$, assuming no evolution of sources. Band et al. (1999) applied the Bayesian methods

to the dataset of Schaeffer (1998) and eight BeppoSAX bursts with good localization and host galaxies. This analysis rules out burst energies below 2×10^{52} ergs, and favors, although not strongly, the value $\approx 10^{53}$ ergs.

3.4. Gravitational lensing searches

Given that GRBs are located at the cosmological distances, some of them should, like all other distant sources, be gravitationally lensed (Paczynski, 1986). Since GRBs are transients, a lensed burst would consist of two (or more) bursts with identical spectra and time profiles, however delayed in time by typically a month. Mao (1992) first estimated of the lensing probability to lie between 0.05% and 0.4%. The lensing probability decreases with the time delay on a timescale of a month, and it was found to peak at $\Delta t \approx 50s(M_*/10^6 M_\odot)$, where M_* is the mass of the lensing object. Thus acquiring a large enough database of bursts should allow to find lensed gamma-ray bursts. However one has to bear in mind that there are some significant selection effects against detecting lensed events. First, BATSE does not monitor all sky simultaneously, the average exposure being about 40% of the sky time. Moreover, after detecting a burst BATSE turn into inactive mode for a 90 minutes. If a burst happens at this time it is only recorded if it is stronger than the preceding burst however, than the information on the preceding burst is lost. Thus detecting lensed bursts with time delays up to 90 minutes is hardly possible. Searches for lensed GRB in the BATSE database were conducted Nemiroff et al. (1994); Marani et al. (1998) yielding null results.

The gravitational lensing problem has recently been revisited by Holz et al. (1999), who found that actually including multiple lensing is important. They have calculated lensing probabilities as a function of the average redshift z_{ave} of the GRB distribution for different cosmologies. Basing on no detection of GRB lensing they derive an upper limit on the average redshift $z_{ave} < 2.2, 2.8, 4.3, 5.3$ for (Ω, Λ) pairs $(0.3, 0.7)$, $(0.5, 0.5)$, $(0.5, 0.0)$, and $(1.0, 0.0)$ respectively. It should be noted that the gravitational lensing probes the spatial distribution of GRBs directly and is not very sensitive to the luminosity function.

3.5. Clustering

While the overall distribution of GRBs shows an amazing degree of isotropy there is also a question of isotropy or clustering on the small angular scales. There has been evidence for such anisotropy - small scale clustering of burst - in the first BATSE catalogue found by Quashnock and Lamb (1993b), who interpreted it as evidence for burst repeating. However, with the change of the BATSE localization algorithm the small scale clustering detection was not confirmed, and has not been seen in the larger data sets.

Within the framework of the cosmological model small scale clustering is expected, if bursts trace luminous matter. Lamb and Quashnock (1993) suggested that given a large enough sample of bursts the large scale structure in the Universe should be detectable: 1000 bursts should suffice to probe the supercluster scales, and about 3000 bursts should probe the scales above $25\,h^{-1}$ Mpc. These results were used with the BATSE 3B catalogue to find that the median distance to the weak bursts must be larger than $630h^{-1}$ Mpc, and the closest burst must be further away than $40h^{-1}$ Mpc (Quashnock, 1996). While these



Figure 5. The left panel shows seven Beppo SAX burst with measured redshifts in the z vs peak flux plane. If GRBs were standard candles the points would lie on a straight line. The right panel shows the cumulative distribution of peak photon luminosities for SAX bursts with redshifts. I have omitted GRB980425. The observed luminosity function is almost a factor of 10^3 wide.

limits are not very constraining they grow approximately linearly with the number of bursts detected provided that no clustering is found. Alternatively, the number of GRBs in the current BATSE database should soon allow to probe the large scale structure.

3.6. Direct measurements

The studies of GRB spatial distribution have entered a new era in the beginning of 1997 with the BeppoSAX discovery of X-ray afterglows (Costa et al., 1997b), This led to rapid optical followups and the discovery of optical afterglows (Groot et al., 1997a), and consequently to identification of host galaxies (Groot et al., 1997b). The accurate optical spectroscopy of either the afterglows themself of the host galaxies allows to find GRB redshifts, with the first redshift of 0.835 measured for GRB970508 (Metzger et al., 1997b). At the time of writing there are nine GRB redshifts measured, including two which are rather uncertain. A list of the GRBs with measured redshifts is presented in Table 2. This table was prepared with the aid of the information posted on Jochen Greiners web page (Greiner, 1999).

We present the "Hubble" diagram for GRBs in the left panel of Figure 5. There is no clear correlation between the redshift and peak flux, I even had to use the logarithmic scale to show all the fluxes. This indicates that the GRB luminosity function is broad. In the right panel of Figure 5 I present the cumulative distribution of peak photon luminosities for Beppo SAX bursts with redshift. In this Figure I have omitted two ASM/RXTE bursts from Table 2 and also GRB980425 which, if the identification with SN1998bw is correct would be five orders of magnitude fainter. The luminosities have been calculated assuming that $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$, and $H_0 = 65\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$. The observed

Table 2. GRBs with measured redshifts.

GRB	z	F_{peak} cts s^{-1}	F_{WFC} [Crab]	Reference
970228	0.695	3700	0.23	Costa et al. (1997a)
970508	0.835	450	1.0	Costa et al. (1997c)
970828	0.3[b]		0.735[a]	Remillard et al. (1997)
971214	3.42	650	1.0	Heise et al. (1997)
980329	5 [c]	6000	6.0	Frontera et al. (1998)
980425	0.008[d]	350	3.0	Soffitta et al. (1998)
980613	1.096	350	0.6	Smith et al. (1998)
980703	0.96		1.7[a]	Levine et al. (1998)
990123	1.60	11200	3.4	Feroci et al. (1999)

[a] RXTE-ASM fluxes between 2 and 12 keV

[b]Redshift of 0.3 is based on a weak detection of iron line (Yoshida, 1998); interpreting the spectral feature as the iron edge the redshift becomes ≈ 0.9.

[c]Redshift determined from broad band photometry (Fruchter, 1999). A faint galaxy at the same location has been found, however it lies at a lower redshift.

[d]Redshift determined on the assumption that SN1998bw is the counterpart of the GRB (Galama et al., 1998).

luminosity function is a factor of almost 10^3 wide, and this result has been obtained already with a small sample of a few burst. This is an indication that the intrinsic luminosity of function of GRBs could be even broader than that!

With the measurements of redshifts we can now directly probe the spatial distribution of GRBs. The cumulative distribution of GRB redshifts measured to date is presented in Figure 6. I have included all the entries from Table 2 in this Figure, despite the fact that, as discussed above, some of the entries could be uncertain. Also since we know that the luminosity function is very broad we may not be detecting all the bursts from high redshifts. Therefore studies of GRBs with more sensitive instruments are very important.

Models for GRBs associate them with late stages of stellar evolution, either compact object mergers, or collapsars. Thus one expects a relation between the star formation rate and the GRB bursting rate. The star formation rate has recently been measured in the Hubble Deep Field (Madau et al., 1996), which shows a dramatic change in the star formation rate between $z = 1$ and present. Totani (1997) calculated the expected GRB rate based on the known star formation history and the synthesis of stellar (and binary) evolution. Wijers et al. (1998) fitted the combined BATSE and PVO brightness distribution with a star formation model, and the dimmest burst could come from the redshifts as high as $z \approx 6$. Based on their results I present a model with the GRB rate proportional to star formation rate in Figure 6. While the two curves do not match very well, they look similar despite the fact that no selection effects due



Figure 6. Cumulative spatial distribution of GRBs from measured redshifts. All entries from Table 2 are shown.We present a model where the GRB rate is proportional to star formation rate (dashed line) (Wijers et al., 1998).

to detector sensitivity, or luminosity function were taken into account. This could be an indication that GRBs are connected with star formation.

4. Classes of GRBs

All that has been written above was based on the assumption that there is only one population of GRBs. There are however some indications that there may exist different classes of GRBs. One of the first BATSE results was to show the bimodality of the duration distribution (Kouveliotou et al., 1993), with typical durations of less than a second and a few tens of seconds. At the same time it has been found that GRBs can be grouped according to their duration and variability properties and form two classes - short variable, and long smooth (Lamb et al., 1993). Moreover brightness distributions of bursts with different hardness ratios are different Pizzichini (1995). The low hardness ratio bursts show a typically



Figure 7. The left panel shows BATSE gamma ray bursts in the duration - hardness plane. Two groups of bursts are clearly visible: the long and the short burst. The long bursts are typically softer than the short ones. The right panel shows the peak flux distributions of the long ($T_{90} > 2s$, the thick line) and the short bursts ($T_{90} < 2s$, the thin line). The two distributions are different, and assuming that the sensitivity to both classes of bursts is similar, the probability that they come from the same distribution is $\approx 10^{-13}$.

steeper behavior in the cumulative brightness distributions, while the harder bursts are those that actually show the rollover in this distribution.

In the left panel Figure 7 I present the BATSE bursts in the duration-hardness plane (hardness is defined as the ratio of the fluxes in BATSE channels 2 and 3). The duration bimodality is clearly seen, moreover the long bursts have typically softer spectra than the short ones, although there is a significant overlap. The properties of GRBs have been analyzed by dividing them into different classes on the hardness duration plot. Belli (1995) used a line defined by $HR = 0.5T_{90}^{1/2}$. Tavani (1998) divides the bursts into four groups by two lines: duration of 2.5 s, and hardness ratio $HR = 3$. Only the long hard bursts $\log(N)$-$\log(S)$ distribution shows deviation from the Euclidean $-3/2$ slope. I present the peak flux distributions of the long and short burst in the right panel Figure 7. Such different brightness distributions could indicate that there are two populations with different luminosities. On the other hand a similar effect has been found when analyzing distribution the peaks in BATSE bursts (Pendleton et al., 1997). Peaks in GRBs evolve typically from hard to soft spectra, and one could speculate that the short bursts are just single peaked, while the longer ones contain more of the soft emission from the tails of the peaks. It should be noted that SAX probes only long bursts (duration larger than 6 s) and so far all but one burst have been in the hard category.

The discovery of an unusual supernova in the errorbox of GRB980425 (Galama et al., 1998) has sparked a debate on a possibility of the association of GRBs and supernovae. If the association were correct then gamma-ray luminosity of GRB980425 would be much smaller than the typical luminosity of

GRBs. It is still unclear whether this association is correct and if there exists a class of supernova related GRBs. If it does then it would be interesting to find out what fraction of GRBs this class constitutes, and whether the supernovae related bursts are related to one of the GRB classes shown above. If there exists a class of under luminous, supernova related GRBs then the faint bursts should be dominated by distant bursts in this class. A future very sensitive experiment may resolve this, however it looks more promising to pursue better quick localizations and multi wavelength followups.

5. Summary

Gamma-ray bursts sky distribution is consistent with isotropy to a very high degree. The estimates of the distance scale and spatial distribution of GRBs seem to converge. The typical gamma-ray burst lie at redshifts of 1 to 2, however this distribution probably has a long tail extending to higher redshifts. GRB luminosity function is broad; from the observation of just a few bursts we already see that the observed luminosity function is almost a factor of one thousand wide. This means that the luminosity function plays a crucial role in shaping the GRB brightness distribution. One of the the brightest bursts seen by BATSE - GRB990123 - lies at the redshift of 1.60! It is possible that there exist different classes of GRB within what we call now cosmic GRBs. This could reflect either different physical mechanism connected with the central engine (compact object mergers or collapsars), or perhaps even orientation effects. Only the long and hard bursts duration distribution exhibits a rollover characteristic for the cosmological models, and this could be an indication that the remaining bursts with Euclidean brightness distributions are different. Finally, one should note that while the evidence for identification of GRB980425 with SN1998bw is still uncertain, it is certainly worth to investigate a possible GRB-supernova connection.

Acknowledgments. The author wishes to thank the organizers for making the Graftavallen meeting a success in both science and fun categories. I would like to thank S. Bajtlik, K. Belczyński, and M. Chodorowski for help in preparing this manuscript. This work has been supported by the KBN grant 2P03D00415.

References

Band D.L., Hartmann D.H., and Schaefer B.E., 1999, ApJ, 514, 862

Belli B.M., 1995, Ap&SS, 231, 43

Bulik T., Lamb D.Q., and Coppi P.S., 1998, ApJ, 505, 666

Cohen E. and Piran T., 1995, ApJ, 444, L25

Costa E., Feroci M., Frontera F., et al., 1997a, IAU Circ, 6572, 1

Costa E., Feroci M., Piro L., et al., 1997b, IAU Circ, 6576, 1

Costa E., Feroci M., Piro L., et al., 1997c, IAU Circ, 6649, 1

Davis S.P., 1995, Ph.D. thesis, The Catholic University of America.

Dermer C.D., 1992, Physical Review Letters, 68, 1799

Fenimore E.E. and Bloom J.S., 1995, ApJ, 453, 25

Fenimore E.E., In'T Zand J.J.M., Norris J.P., Bonnell J.T., and Nemiroff R.J., 1995, ApJ, 448, L101

Feroci M., Piro L., Frontera F., et al., 1999, IAU Circ, 7095, 1

Fishman G.J., Meegan C.A., Wilson R.B., et al., 1994, ApJS, 92, 229

Frontera F., Costa E., Piro L., et al., 1998, IAU Circ, 6853, 1

Fruchter A.S., 1999, ApJ, 512, L1

Galama T.J., Vreeswijk P.M., Van Paradijs J., et al., 1998, Nature, 395, 670

Greiner J., 1999, `http://www.obs.apm.de/~jcg/grbgen.html`

Groot P.J., Galama T.J., van Paradijs J., et al., 1997a, IAU Circ, 6584, 1

Groot P.J., Galama T.J., Van Paradijs J., et al., 1997b, IAU Circ, 6588, 1

Hakkila J., Meegan C.A., Horack J.M., et al., 1996, ApJ, 462, 125

Heise J., In 'T Zand J., Spoliti G., et al., 1997, IAU Circ, 6787, 1

Holz D.E., Miller M.C., and Quashnock J.M., 1999, ApJ, 510, 54

Horack J.M., Emslie A.G., and Meegan C.A., 1994, ApJ, 426, L5

Horvath I., Meszaros P., and Meszaros A., 1996, ApJ, 470, 56

Klebesadel R., Strong I., and Olson R., 1973, ApJ, 182, L85

Kouveliotou C., Meegan C.A., Fishman G.J., et al., 1993, ApJ, 413, L101

Lamb D.Q., 1995, PASP, 107, 1152

Lamb D.Q., Graziani C., and Smith I.A., 1993, ApJ, 413, L11

Lamb D.Q. and Quashnock J.M., 1993, ApJ, 415, L1

Larson S.B. and MCLean I.S., 1997, ApJ, 491, 93

Lee T.T. and Petrosian V., 1997, ApJ, 474, 37

Levine A., Morgan E., and Muno M., 1998, IAU Circ, 6966, 1

Loredo T.J. and Wasserman I.M., 1995, ApJS, 96, 261

Loredo T.J. and Wasserman I.M., 1998a, ApJ, 502, 75

Loredo T.J. and Wasserman I.M., 1998b, ApJ, 502, 108

Madau P., Ferguson H.C., Dickinson M.E., et al., 1996, MNRAS, 283, 1388

Mao S., 1992, ApJ, 389, L41

Mao S. and Paczynski B., 1992, ApJ, 388, L45

Marani G.F., Nemiroff R.J., Norris J.P., Hurley K.C., and Bonnell J.T., 1998, in *Huntsville Gamma-Ray Burst Symposium*, 166

Meegan C.A., Paciesas W.S., Pendleton G.N., et al., 1998, in *Huntsville Gamma-Ray Burst Symposium*, 3

Meszaros A. and Meszaros P., 1996, ApJ, 466, 29

Metzger M.R., Cohen J.G., Chaffee F.H., , and Blandford R.D., 1997a, IAU Circ, 6676, 3

Metzger M.R., Djorgovski S.G., Kulkarni S.R., et al., 1997b, Nature, 387, 878

Mitrofanov I.G., Chernenko A.M., Pozanenko A.S., et al., 1996, ApJ, 459, 570

Nemiroff R.J., Wickramasinghe W.A.D.T., Norris J.P., et al., 1994, ApJ, 432, 478

Norris J.P., Bonnell J.T., Nemiroff R.J., et al., 1995, ApJ, 439, 542

Norris J.P., Nemiroff R.J., Scargle J.D., et al., 1994, ApJ, 424, 540

Pacisas W.S., Meegan, C.A., Pendleton, G.F., 1999, ApJS, in press, astro-ph/9903205

Paczynski B., 1986, ApJ, 308, L43

Paczynski B., 1995, PASP, 107, 1167

Pendleton G.N., Paciesas W.S., Briggs M.S., et al., 1997, ApJ, 489, 175

Piran T., 1992, ApJ, 389, L45

Pizzichini G., 1995, in *Porceedings of the 24th International Cosmic Ray Conference in Rome*, 81

Quashnock J.M., 1996, ApJ, 461, L69

Quashnock J.M. and Lamb D.Q., 1993a, MNRAS, 265, L45

Quashnock J.M. and Lamb D.Q., 1993b, MNRAS, 265, L59

Remillard R., Wood A., Smith D., and Levine A., 1997, IAU Circ, 6726, 1

Schaeffer B., 1992, in *Gamma Ray Bursts: Observations, Analyses, Theories*, Cambridge University Press

Schaeffer B., 1998, astro-ph/9807069

Smith M.J.S., Ricci R., In 'T Zand J., et al., 1998, IAU Circ, 6938, 1

Soffitta P., Feroci M., Piro L., et al., 1998, IAU Circ, 6884, 1

Stern B., Poutanen J., and Svensson R., 1997, ApJ, 489, L41

Stern B.E., 1996, ApJ, 464, L111

Tavani M., 1998, ApJ, 497, L21

Totani T., 1997, ApJ, 486, L71

Ulmer A. and Wijers R.A.M.J., 1995, ApJ, 439, 303

Ulmer A., Wijers R.A.M.J., and Fenimore E.E., 1995, ApJ, 440, L9

Wickramasinghe W.A.D.T., Nemiroff R.J., Norris J.P., et al., 1993, ApJ, 411, L55

Wijers R.A.M.J., Bloom J.S., Bagla J.S., and Natarajan P., 1998, MNRAS, 294, L13

Woods E. and Loeb A., 1995, ApJ, 453, 583

Yoshida A., 1998, Talk at the Rome GRB Meeting